

Twistors versus harmonics

C. Lovelace
Physics Dept., Rutgers University,
Piscataway, NJ 08854-8019.

lovelace @ physics.rutgers.edu

June 21, 2010

Abstract

By lifting full Yang-Mills theory to $R_4 \times S_2$, Mason et al. obtained MHV vertices by gauge transformation. Their Lagrangian depended on long intricate twistor manipulations. Spinor S_2 harmonics give a one page proof, eliminating all the CP_3 apparatus.

1 Introduction

It is fashionable to abandon Lagrangians completely and construct twistor S matrices. However survivors from the 1960's have unpleasant memories of a similar fashion. Still interesting therefore are a series of five papers by Mason et al. [1, 2, 3]. They claimed that full Yang-Mills [YM] theory can be lifted to $R_4 \times S_2$, and then GAUGE transformed to an MHV basis. This ought to be much simpler than Mansfield's canonical transformation [4]. Unfortunately they wrote in a hieratic dialect (twish?) dear to Oxford mathematicians. It is important to find a similar lifted Lagrangian for supergravity where previous attempts have foundered [5], so a demotic translation would be useful. I will only consider bosonic YM in Euclidean spacetime. Fermions would just complicate the notation.

In two dimensions free wave equations can be solved just by going to complex (or lightcone) coordinates. In 4D this is ambiguous – there are S_2 possibilities and there is no reason to make the same choice at each point. Thus one is led to consider $R_4 \times S_2$ in order to make analytic or lightcone gauge covariant [6]. Twistor people regard $R_4 \times S_2 = C_2 \times CP_1$ as a stepping stone to CP_3 , but I will not pursue that. Instead of twistors I will use the S_2 harmonics of Galperin et al. [7], which are a spinor extension of familiar spherical harmonics. They were used long ago [8] to solve selfdual YM. I will first review this (§2,3), and then extend it to full YM theory (§4). In §5 I check that the result is equivalent to Mason's. After explaining prerequisites, my proof fits into one page.

2 Spinor harmonics

Here x is a point in Euclidean spacetime; u, v, w are points on the fiber S_2 . Since Lagrangians are local on the R_4 base, no fiber bundle theory is needed. D denotes an $su(2)$ generator on the fiber. Otherwise partial derivatives are ∂ . When gauge potentials A are added, I call both types ∇ . The $\pm$ superscripts count a conserved charge on the S_2 fiber.

In standard spinor notation [9], irreducible representations of $Spin(4) = SU(2)_L \times SU(2)_R$ are $\psi_{(AB\dots)(A'B'\dots)}$, where $(\)$ means symmetrize. Indices are raised or lowered by $\psi^A = \epsilon^{AB}\psi_B$, $\psi_A = \psi^B\epsilon_{BA}$, where $\epsilon_{AB} = \epsilon^{AB} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$.

Consider $su(2)_R$. We could represent it on S_2 by the usual angular momentum operators $L_{\pm}, L_0$ expressed in polar coordinates θ, ϕ . However as shown by Galperin et al.[7], it is more convenient to label a point $u \in S_2$ by spinors $u^{+A'}$, $u_{B'}^-$, satisfying (in the above sign convention)

$$u^{+A'} u_{A'}^- = 1 = -\epsilon^{B'A'} u_{B'}^+ u_{A'}^-, \quad (1)$$

$$(u^{+A'})^* = -u_{A'}^-, \quad (2)$$

where $*$ means complex conjugate. Eqs.(1)-(2) could be solved by a single complex number u_1^+ , as in stereographic projection, but they are more powerful unsolved (to avoid patching). The $su(2)_R$ generators are then

$$D^{\pm\pm} \equiv L_{\pm} = u^{\pm A'} \frac{\partial}{\partial u^{\mp A'}}, \quad D^0 \equiv 2L_0 = u^{+A'} \frac{\partial}{\partial u^{+A'}} - u^{-A'} \frac{\partial}{\partial u^{-A'}}. \quad (3)$$

Since $S_2 = SO(3)/SO(2)$, the D^0 eigenvalue is required to be a conserved charge. Its value is the difference between the number of + and - superscripts. (This reduces $su(2)$ representations to the coset space.) If u, v are two points on S_2 , define

$$\langle uv \rangle \equiv \epsilon^{A'B'} u_{A'}^+ v_{B'}^+. \quad (4)$$

$\langle uv \rangle$ has charge $++$, but I will only show this when it matters. An important identity [7] is

$$\langle u | (D^{++})^{-1} | v \rangle = \langle uv \rangle^{-1}. \quad (5)$$

This is the bilocal Green's function on S_2 that inverts the differential operator $D^{++} \equiv L_+$ (subject of course to global conditions).

Euclidean spacetime coordinates x^μ can be converted to bispinors by

$$x^{BB'} = ix^4 + (\vec{\sigma} \cdot \vec{x}), \quad (6)$$

and then shuffled into analytic (+) and anti-analytic (-) doublets

$$y^{\pm B} \equiv u_{B'}^{\pm} x^{BB'}. \quad (7)$$

Similarly

$$\partial_B^{\pm} = u^{\pm B'} \partial_{BB'}. \quad (8)$$

Note that by (1) the signs reverse

$$\partial_B^\pm = \pm \partial / \partial y^{\mp B}. \quad (9)$$

Now add gauge potentials (antihermitian matrices)

$$\nabla_{BB'} \equiv \partial_{BB'} + A_{BB'}. \quad (10)$$

It is well known [9] that

$$[\nabla_{BB'}, \nabla_{CC'}] = F_{(BC)} \epsilon_{B'C'} + F_{(B'C')} \epsilon_{BC}. \quad (11)$$

This separates the field strength into selfdual $F_{(BC)}$ and antiselfdual [ASD] components. Thus far x, u are independent. Multiplying by harmonics

$$\nabla_B^\pm \equiv u^{\pm B'} \nabla_{BB'}, \quad (12)$$

(11) gives

$$[\nabla_B^+, \nabla_C^+] = \epsilon_{BC} u^{+B'} u^{+C'} F_{(B'C')}, \quad (13)$$

an equation which will be very useful later.

3 Selfdual Yang-Mills

Now suppose the YM theory is selfdual $F_{(B'C')} = 0$. Here I am summarizing [8]. Then

$$[\nabla_B^+, \nabla_C^+] = 0. \quad (14)$$

So by Frobenius' theorem [10] there exists a gauge transformation $\Lambda(x, u)$ that locally flattens this connection. Since it depends on u , it will unflatten the $su(2)_R$ algebra (3). Thus OLD to NEW gauge transforms

$$\nabla_B^+ \rightarrow \partial_B^+, \quad D^{\pm\pm} \rightarrow \nabla^{\pm\pm} \equiv D^{\pm\pm} + A^{\pm\pm}. \quad (15)$$

The two new components $A^{\pm\pm}$ are enough to compensate for the two old components A_B^+ , so D^0 can stay flat.

By (3) and (12) in old gauge

$$[D^{++}, \nabla_B^+] = 0, \quad (16)$$

so in new gauge

$$[\partial_B^+, \nabla^{++}] = 0, \quad (17)$$

which means by (9) that A^{++} is independent of y_B^- . The $su(2)_R$ algebra becomes in new gauge

$$[\nabla^{++}, \nabla^{--}] = D^0, \quad (18)$$

with no potential on the rhs. As explained in [8], this can be used to solve for A^{--} in terms of A^{++} . The result is most easily described by perturbing the Green's function (5)

$$\begin{aligned} \mathcal{G}^{--}(u, v) &\equiv \langle u | [D^{++} + A^{++}]^{-1} | v \rangle \\ &= \langle uv \rangle^{-1} - \int dw \langle uw \rangle^{-1} A^{++}(x, w) \langle wv \rangle^{-1} + \dots \end{aligned} \quad (19)$$

The n^{th} term of the perturbation expansion has $n - 1$ integrals over S_2 , while x stays fixed. Then by [8] eq.(III.26),

$$- A^{--}(x, u) = \lim_{v \rightarrow u} [\mathcal{G}^{--}(u, v) - \langle uv \rangle^{-1}] . \quad (20)$$

Because of the singular denominators this *does* depend on y_B^- .

Lastly we can find the nonzero part of the field strength $F_{(BC)}$. In old gauge, (3) and (12) gave

$$\nabla_B^- = [D^{--}, \nabla_B^+] . \quad (21)$$

In new gauge this becomes by (15)

$$A_B^- = -\partial_B^+ A^{--} . \quad (22)$$

In old gauge, (11) with $F_{(B'C')} = 0$ gave by (1) and (12)

$$[\nabla_B^+, \nabla_C^-] = -F_{(BC)} . \quad (23)$$

Thus in new gauge

$$F_{(BC)} = \partial_C^+ \partial_B^+ A^{--} . \quad (24)$$

So the selfdual theory is entirely determined by the unconstrained prepotential $A^{++}(y_B^+, u)$, which is independent of y_B^- by (17). Of course similar results were obtained earlier by Ward [10] using twistors, but they are not as easy to follow as [8].

4 Full Yang-Mills

Let us see how rapidly this technique reproduces long esoteric twistor manipulations. For completeness I will first put the usual YM Lagrangian into Chalmers-Siegel form.

Let $F^\pm$ be the SD/ASD YM field strengths as in (11). It is well known [6, 11] that $(F^+)^2 - (F^-)^2$ is a total divergence. Therefore the standard YM Lagrangian (trace over gauge group assumed)

$$\mathcal{L} = \frac{1}{16} F^{\mu\nu} F_{\mu\nu} = \frac{1}{8} [(F^+)^2 + (F^-)^2] \quad (25)$$

is equivalent to

$$\mathcal{L} = \frac{1}{4} (F^-)^2 \equiv \frac{1}{4} F^{(C'D')} F_{(C'D')} . \quad (26)$$

It is convenient to introduce a dummy field $G^{(C'D')}(x)$, which can be functionally integrated out by completing the square, leaving (26):

$$\mathcal{L} = \frac{1}{2} G^{(C'D')} F_{(C'D')} - \frac{1}{4} G^{(C'D')} G_{(C'D')} . \quad (27)$$

If we omitted the second term, G would be a Lagrange multiplier for selfdual YM [12].

Now the race starts. By (13)

$$[\nabla_C^+, \nabla_D^+] = \epsilon_{CD} u^{+C'} u^{+D'} F_{(C'D')} . \quad (28)$$

This suggests a natural way to include $G^{(C'D')}$. Introduce a scalar potential $B^{--}(x, u)$ and identify $G^{(C'D')}$ with three of its Fourier modes on S_2

$$G^{(C'D')}(x) = \int du u^{+C'} u^{+D'} B^{--}(x, u) . \quad (29)$$

Then the first term of (27) becomes

$$\frac{1}{2} G^{(C'D')} F_{(C'D')} = \frac{1}{4} \int du B^{--}(x, u) \epsilon^{CD} [\nabla_C^+, \nabla_D^+] . \quad (30)$$

At first sight the second term of (27) becomes

$$-\frac{1}{4} G^{(C'D')} G_{(C'D')} = -\frac{1}{4} \int du \int dv B^{--}(x, u) \langle u^+ v^+ \rangle^2 B^{--}(x, v) , \quad (31)$$

where $\langle u^+ v^+ \rangle$ is (4). However this is not gauge invariant. The gauge potentials $B(u)$, $B(v)$ need to be connected by Wilson lines on S_2

$$\exp\left(\int_u^v dw A^{++}(w)\right) ,$$

forming a loop when traced. (In 2D we can fix $A^{--} = 0$.) This is equivalent to inserting propagators (19), but then we need two more $\langle u^+ v^+ \rangle$ factors to conserve the D^0 charge. So the unique correct formula is

$$-\frac{1}{4} \int du \int dv \langle u^+ v^+ \rangle^4 \text{Tr}\{B^{--}(x, u) \mathcal{G}^{--}(u, v) B^{--}(x, v) \mathcal{G}^{--}(v, u)\} , \quad (32)$$

where

$$\mathcal{G}^{--}(u, v) \equiv \langle u | [D^{++} + A^{++}]^{-1} | v \rangle . \quad (33)$$

Check: if $A^{++} = 0$, (32) = (31) by (5). So far A^{++} belongs to a trivial 2D gauge theory independent of x . We can also add a Lagrange multiplier term to implement (16) :

$$B^{---C} [\nabla^{++}, \nabla_C^+] . \quad (34)$$

Now perform a gauge transformation $\Lambda(x, u)$. This will change

$$A^{++}(u) \rightarrow A^{++}(x, u) ,$$

but of course it won't eliminate A_C^+ completely, unlike the SD case (15). We can however set one component $A_2^+ = 0$, thus defining a new gauge. The final Lagrangian is

$$\begin{aligned} & \frac{1}{4} \epsilon^{CD} \int du \text{Tr}\{B^{--} [\nabla_C^+, \nabla_D^+] \} + \int du \text{Tr}\{B^{---C} [\nabla^{++}, \nabla_C^+] \} \\ & - \frac{1}{4} \int du \int dv < uv >^4 \text{Tr}\{B^{--}(x, u) \mathcal{G}^{--}(u, v) B^{--}(x, v) \mathcal{G}^{--}(v, u)\}. \end{aligned} \quad (35)$$

In the gauge $\partial_C^- A^{++} = 0$ it reduces to (27) + (29). In the gauge $A_2^+ = 0$ the unwanted BAA vertex in the first term vanishes, and (19) gives a series of MHV vertices instead. The bilocal S_2 term first appeared in [13] .

5 Comparison

The clearest of the five Mason et al. papers is Ref.[2], where [1] are usefully summarized. (My notations were chosen for easy comparison). Boels starts with the usual twistor equation

$$\omega^C = x^{CC'} \pi_{C'} . \quad (36)$$

Conjugation in Euclidean spacetime is

$$\hat{\pi}_1 = -\bar{\pi}_2 , \quad \hat{\pi}_2 = +\bar{\pi}_1 . \quad (37)$$

Twistors are defined only up to a scale. If we fix this by $< \pi \hat{\pi} > = 1$, we can identify

$$\pi_{C'} = u_{C'}^+ , \quad \hat{\pi}_{C'} = -u_{C'}^- , \quad (38)$$

to get (1)-(2) above. Then $\omega^C = y^{+C}$ by (7). Next, eq.(5) of Ref.[2] becomes by (3) and (9) above

$$\bar{\partial}_0 = D^{++} , \quad \bar{\partial}_\alpha = \partial_C^+ . \quad (39)$$

Their action (eq.(16) of Ref.[2]) is

$$\begin{aligned} S = & \frac{1}{2} \int d^4 x dk B_0 (\bar{\partial}^\alpha A_\alpha + g A^\alpha A_\alpha) + B^\alpha (\bar{\partial}_\beta A_0 - \bar{\partial}_0 A_\beta + g[A_\beta, A_0]) \\ & - \frac{1}{4} \int d^4 x dk_1 dk_2 H_1^{-1} B^0(\pi_1) H_1 H_2^{-1} B^0(\pi_2) H_2 < \pi_1 \pi_2 >^4 . \end{aligned} \quad (40)$$

It can now be matched to (35) above. Clearly $A_0 = A^{++}$, $B_0 = B^{--}$ and the first two terms agree. The third term has the same general structure, but what is H ? “ H is a holomorphic frame of the gauge bundle over $p^{-1}(x)$ such that the covariant derivative of it vanishes on the sphere.” I can’t translate this hieroglyph (it looks upside down), but if we identify

$$H_1 H_2^{-1} = \mathcal{G}^{--}(u, v) , \quad (41)$$

we get agreement since $< \pi_1 \pi_2 > = < uv >$. Eq.(2.49) of Ref.[3], with notation adjusted to match [2], gives their expansion in MHV vertices

$$G^{(C'D')}G_{(C'D')} = - \sum_{n=2}^{\infty} (-i\sqrt{2}g)^{n-2} \int \sum_{p=2}^n \text{tr}(B_0^1 A_0^2 \cdots A_0^{p-1} B_0^p A_0^{p+1} \cdots A_0^p) \\ \times \langle \pi_1 \pi_p \rangle^4 / (\langle \pi_1 \pi_2 \rangle \langle \pi_2 \pi_3 \rangle \cdots \langle \pi_n \pi_1 \rangle), \quad (42)$$

which is exactly what results when (19) is substituted into the last term of (35).

Thus harmonics are much faster and clearer than twistors. The trick is to start with distinct gauge theories on R_4 and S_2 , and then mix them by a gauge transformation. The CP_3 apparatus just confused the issue. Now that a simple intuitive proof of Mason's Lagrangian exists, I hope to extend it to gravity using *local* twistors (heresy in Oxford). The powerful formalism of Ref.[7] could be generalized to any symmetric space, and might have more applications.

References

- [1] L.Mason, JHEP 0510:009 (2005), hep-th/0507269;
R.Boels, L.Mason, and D.Skinner, JHEP 0702:014 (2007), hep-th/0604040;
Phys.Lett.B648,90 (2007), hep-th/0702035.
- [2] R.Boels, Phys.Rev.D76,105027 (2007), hep-th/0703080.
- [3] R.Boels and C.Schwinn, JHEP 0807:007 (2008), arXiv:0805.1197.
- [4] P.Mansfield, JHEP 0603:037 (2006), hep-th/0511264;
J.H.Ettle and T.R.Morris, JHEP 0608:003 (2006), hep-th/0605121.
- [5] V.P.Nair, Phys.Rev.D78,041501(R) (2008), arXiv:0710.4961.
- [6] M.F.Atiyah, *Geometry of Yang-Mills fields*, Pisa 1979.
- [7] A.S.Galperin, E.A.Ivanov, V.I.Ogievetsky, and E.S.Sokatchev,
Harmonic superspace, Cambridge 2001, Ch.4.
- [8] A.S.Galperin, E.A.Ivanov, V.I.Ogievetsky, and E.S.Sokatchev,
Ann.Phys.(NY) 185,1 (1988), §III.
- [9] R.Penrose and W.Rindler, *Spinors and spacetime*, Cambridge 1984.
- [10] R.S.Ward and R.O.Wells, *Twistor geometry and field theory*,
Cambridge 1990, pp.142-3.
- [11] T.Eguchi, P.B.Gilkey, and A.J.Hanson, Phys.Reports 66,213 (1980).
- [12] G.Chalmers and W.Siegel, Phys.Rev.D54,7628 (1996), hep-th/9606061.
- [13] V.P.Nair, Phys.Lett. B214,215 (1988).